

SECRETION OF M. TUBERCULOSIS 30 kDa MAJOR EXTRACELLULAR PROTEIN BY BCG






FIG. 1a **FIG. 1b**


- 1, COOMASSIE STAINED PROTEINS AFTER 4 WEEKS OF GROWTH IN BROTH CULTURE
- 2, IMMUNOBLOT WITH ANTI-30 kDa PROTEIN IgG AFTER 4 WEEKS OF GROWTH IN BROTH CULTURE
- 3, IMMUNOBLOT WITH ANTI-30 kDa PROTEIN IgG AFTER 12 WEEKS OF GROWTH IN BROTH CULTURE WITHOUT HYGROMYCIN

2/13


FIG. 2a*Experiment 1: Skin tests to Protein r30***FIG. 2b***Experiment 2: Skin tests to Protein r30*

3/13



FIG. 3a



Experiment 1: Weights of Guinea Pigs After Challenge



4/13

FIG. 3b*Experiment 2: Weights of Guinea Pigs After Challenge*

5/13
Experiments 1 and 2: Lung CFU




FIG. 4a

Experiments 1 and 2: Spleen CFU




FIG. 4b

Skin Test Response to r30




FIG. 5

6/13






FIG. 6

**FIG. 8**

8/13





FIG. 9


Two-Plasmid System for the Expression of *M. tuberculosis* 30 kDa Protein in *M. bovis* BCG Tice



C = Coomassie-stained SDS-PAGE Gels of Culture Filtrates
I = Immunoblots of Culture Filtrates

FIG. 11

FIG. 12



12/13




FIG. 13

FIG. 14

